



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2017

---

**Reply to: comparing C-MAC videolaryngoscope with direct laryngoscopy for  
emergency intubation**

Ruetzler, Kurt ; Sulser, Simon ; Uebmann, Dirk ; Spahn, Donat R

DOI: <https://doi.org/10.1097/EJA.0000000000000695>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-143919>

Journal Article

Published Version

Originally published at:

Ruetzler, Kurt; Sulser, Simon; Uebmann, Dirk; Spahn, Donat R (2017). Reply to: comparing C-MAC videolaryngoscope with direct laryngoscopy for emergency intubation. *European Journal of Anaesthesiology*, 34(11):786.

DOI: <https://doi.org/10.1097/EJA.0000000000000695>

- 2 Niforopoulou P, Pantazopoulos I, Demestiha T, *et al.* Video-laryngoscopes in the adult airway management: a topical review of the literature. *Acta Anaesthesiol Scand* 2010; **54**:1050–1061.
- 3 Kelly FE, Cook TM. Seeing is believing: getting the best out of videolaryngoscopy. *Br J Anaesth* 2016; **117** (Suppl 1):9–13.
- 4 Sakles JC, Mosier JM, Patanwala AE, *et al.* The utility of the C-MAC as a direct laryngoscope for intubation in the emergency department. *J Emerg Med* 2016; **51**:349–357.
- 5 Levitan RM, Heitz JW, Sweeney M, *et al.* The Complexities of tracheal intubation with direct laryngoscopy and alternative intubation devices. *Ann Emerg Med* 2011; **57**:240–207.
- 6 Hossfeld B, Frey K, Doerges V, *et al.* Improvement in glottic visualisation by using the C-MAC PM video laryngoscope as a first-line device for out-of-hospital emergency tracheal intubation: an observational study. *Eur J Anaesthesiol* 2015; **32**:425–431.
- 7 Sakles JC, Mosier J, Chiu S, *et al.* A comparison of the C-MAC video laryngoscope to the Macintosh direct laryngoscope for intubation in the emergency department. *Ann Emerg Med* 2012; **60**:739–748.
- 8 Vassiliadis J, Tzannes A, Hitos K, *et al.* Comparison of the C-MAC video laryngoscope with direct Macintosh laryngoscopy in the emergency department. *Emerg Med Australas* 2015; **27**:119–125.
- 9 Sakles JC, Mosier JM, Patanwala AE, *et al.* The C-MAC (video laryngoscope) is superior to the direct laryngoscope for the rescue of failed first-attempt intubations in the emergency department. *J Emerg Med* 2015; **48**:280–286.
- 10 Natt BS, Malo J, Hypes CD, *et al.* Strategies to improve first attempt success at intubation in critically ill patients. *Br J Anaesth* 2016; **117** (Suppl 1):i60–i68.

DOI:10.1097/EJA.0000000000000694

## Reply to: comparing C-MAC videolaryngoscope with direct laryngoscopy for emergency intubation

Kurt Ruetzler, Simon Sulser, Dirk Uebmann and Donat R. Spahn

From the Institute of Anaesthesiology (KR, SS, DU, DRS), University Hospital Zürich, Zürich, Switzerland and Departments of Outcomes Research and General Anesthesiology, Cleveland Clinic, Cleveland, Ohio, USA (KR)

Correspondence to Kurt Ruetzler, MD, Departments of Outcomes Research and General Anesthesiology, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, USA

Tel: +1 216 407 4108, fax: +1 216 444 2294; e-mail: kurt.ruetzler@reflex.at

Editor,

We appreciate the interest of Liu *et al.*<sup>1</sup> in our publication<sup>2</sup>. We would like to comment on the questions raised in their letter.<sup>1</sup>

First, all three anaesthetists were properly trained in both techniques (C-MAC videolaryngoscope and direct laryngoscopy, University Hospital Zurich, Zurich, Switzerland) prior to the start of this study. Of course, all three had much greater experience with direct laryngoscopy (average about 1000 intubations) compared with C-MAC (between 30 and 300 intubations), but their level

of training was sufficient. The sufficient level of training was confirmed by the finding of our study that the success rate at the first intubation attempt was 100 and 98%, respectively. Second, we basically agree that switching the intubation technique from video to direct laryngoscopy might be an important advantage of several videolaryngoscopes. However, this was neither investigated, nor indicated, in any intubation attempt to achieve successful intubation in our study. Although the clinical assessment of Liu *et al.* may be right, scientific evidence for their conclusion is incomplete. Third, at our institution, a stylet is mandatorily used for all rapid sequence intubation procedures. Therefore, all (endotracheal tubes) in both groups were prepared with a hockey-stick shaped stylet in advance.<sup>3</sup> Fourth, we agree with Liu *et al.* that a cut-off time might be important in some airway studies. However, in the clinical and especially the emergency setting, clinical judgement and decision-making is more important than any definition of time-frame. In general, the rate of successful endotracheal intubation and avoidance of unrecognised oesophageal intubation are clinically of the highest importance. In contrast, the clinical impact of time to intubation is questionable as a difference of a couple of seconds is usually of no clinical relevance. Finally, the well tolerated use of the videolaryngoscope is completely supported by the data of our study and we, therefore, basically agree that the use of the videolaryngoscope might be an acceptable first-line intubation device. However, any recommendation for the first-line device during emergency airway management should be based on the provider's experience and the availability of the device.

## Acknowledgements related to this article

Assistance with the reply: none.

Financial support and sponsorship: none.

Conflicts of interest: none.

## References

- 1 Liu Y-Y, Xue F-S, Li H-X, Yang G-Z. Comparing C-MAC videolaryngoscope with direct laryngoscopy for emergency intubation. *Eur J Anaesthesiol* 2017; **34**:785–786.
- 2 Sulser S, Uebmann D, Schlaepfer M, *et al.* C-MAC videolaryngoscope compared with direct laryngoscopy for rapid sequence intubation in an emergency department: a randomised clinical trial. *Eur J Anaesthesiol* 2016; **33**:943–948.
- 3 Sulser S, Uebmann D, Brueesch M, *et al.* The C-MAC videolaryngoscope compared with conventional laryngoscopy for rapid sequence intubation at the emergency department: study protocol. *Scand J Trauma Resusc Emerg Med* 2015; **23**:38.

DOI:10.1097/EJA.0000000000000695